

Air quality in megacities Faraday Discussion

Concluding Remarks Lecture
Nov 20, 2020

Chak K. Chan (City University of Hong Kong, China)

Masao Gen (Kanazawa University, Japan)




Theme 1

Current status and trends in air quality in megacities

PM2.5 (Guangzhong Basin, SJV, Beijing, LA Basin, Shanghai)

- Importance of BC, OC, ammonium (Watson et al., GZB vs. SJV)
- OC measurements and comparison of source apportionment techniques of PM in Beijing CMB, EG (C14), ACSM/AMS PMF, filter based PMF (Xu et al.)
- Real time tracer-based OC speciation by TAG (Zhu et al., Shanghai)
- Importance of vehicular emission controls, tailpipe and non-tailpipe (Altuwayjiri et al., LA Basin)

Absolute contribution of both tailpipe emissions and non-tailpipe emissions to OC decreased in LA Basin



- Tailpipe emissions reduced drastically.
- Non-tailpipe emissions was replacing tailpipe emissions to be the primary source.
- Regulations reduced the emissions of primary organic precursors ——>SOA↓

Altuwayjiri et al., Long-term trends in the contribution of PM2.5 sources to organic carbon (OC) in the Los Angeles basin and the effect of PM emission regulations


VOC, Ozone and Ox (London, Bangkok, Delhi, Asia)

- Decreasing trend of NOx in both London and Bangkok roadside stations in 2005-2015, but less obvious reductions in O₃ (Khan et al.)
- Despite widespread open biomass burning in India, traffic related petrol and diesel emissions remain the key drivers of gas-phase urban air pollution in Delhi (Stewart et al.)
- VOC limited Ozone formation in Delhi (Chen et al.)
 - 50% reduction in AOD increases maximum hourly ozone concentrations by 5% in June, but increases it by 25% in November.
- Asia road emission trends Emissions of NOx, CO, and PM2.5 peaked in 2000-2015 but GHG will only peak in 2040 (He et al.)

COVID 19 – lockdown

- Reduction of NOx, SOx, CO in Beijing; weekly and diurnal patterns become less obvious. (Brimblecombe and Lai)
- Tracer based work suggests increase in secondary PM, especially sulfate and SOA. (Zhu et al.)
- Other literature has also reported the increase of ozone and secondary PM2.5 in mega cities in China during COVID 19

Difference of $PM_{2.5}$ contribution between BR and DR by (a) individual secondary factors and (b) individual primary factors.



- Contribution of primary source factors to PM_{2.5} and OC dropped (highest contributors: vehicular emissions, biomass burning, coal combustion)
- Contribution of secondary sources factors (except nitrate) to PM_{2.5} and OC increased; mainly due to sulfate formation and SOA related to later generation organic oxidation products


Zhu et al. Tracer-based Characterization of Source Variations of PM2.5 and Organic Carbon in Shanghai Influenced by the COVID-19 Lockdown

Theme 2

Physico-chemical processes in the urban atmosphere (neighbourhood scale)


Impact of secondary aerosol formation on PM_{2.5} concentrations over street network in Paris (Lugon et al.)

 $PM_{2.5}$ simulated with and without processes leading to secondary aerosol formation (gas chemistry and aerosol dynamics)



- Necessity to consider secondary aerosol formation in streets
- Higher impacts (>18%) of secondary aerosol formation on PM_{2.5} concentrations were observed in streets with high traffic strength

Plume characteristics in a cluster of tall buildings (TB) (Hertwig et al.)



Aerosol-boundary layer feedback via coupled LES aerosol-radiation model (Slater et al.)





Fig. 6 : Aerosol concentration vs PBL % decrease for low initial PBL simulations and original PBL simulations, due to different aerosol loadings

Policy implication : Control of BC


Ozonolysis of cooking emissions (Milsome et al.)





- Kitchen emissions in Hong Kong
- Traffic emissions in Gothenburg
- Primary and Secondary Aerosols


Roadside campaign in Mong Kok (URBAN) March – July, 2013; AMS + others














13

American Geophysical Union (AGU)



reactions of particles in the sun add organic aerosols to the atmosphere. These tiny particles are a major component of airborne. particle pollution and can cause heart and lung problems in humans and reduce visibility, according to the U.S. Environmental Protection Agency.

Hong Kong, a city of more than 7 million people, has struggled with air pollution. At times, much of the city's famous skyline along the coast is masked by a thick layer of smog, according to the study's authors.

The new study sought to identify the sources of organic aerosols in Hong Kong by continuously monitoring fine particles on an urban street in the Mong Kok area of downtown Hong Kong.

The study found that from March to July 2013, cooking aerosols accounted for 35 percent of organic aerosols in the air, while 25 percent of the organic aerosols came from traffic. The remaining 40




Berto Lee (left) and Chak Chan (right) stand in a shelter in Hong Kong's Mong Kok next to the Aerodyne High Resolution Aerosol Mass Spectrometer, used to sample airborne particulate matter.



Credit: Chak Chan

Ideas and opinions expressed on this site are those of the authors and commenters alone. They do not necessarily represent the views of the American Geophysical Union.

OTHER AGU BLOGS














Cooking: important source of OA



(Allan et al., 2010; Sun et al., 2011; 2013; Mohr et al., 2012; Crippa et al., 2013; Lee et al., 2015)

SOA formation from gas-phase emissions of heated cooking oils






Alkanals and alkenals, dominated the non-methane organic gas (NMOG) emissions

SOA: one order of magnitude higher than POA

PR=[SOA]×Dilution ratio×Flow rate ER=[POA]×Dilution ratio×Flow rate










Roadside Assessment of a **Modern City Bus Fleet**: Primary and Secondary Emissions



Qianyun Liu¹, Åsa M. Hallquist², Henrik Fallgren², Martin Jerksjö², Sara Jutterström², Håkan Salberg², Mattias Hallquist³, Michael Le Breton³, Xiangyu Pei³, Ravi Kant Pathak³, Tengyu Liu⁴, Berto Lee⁴, and Chak K. Chan⁴


¹Division of Environment and Sustainability, Hong Kong University of Science and Technology, China

²IVL Swedish Environmental Research Institute, Gothenburg, Sweden

³Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg, Sweden

⁴School of Energy and Environment, City University of Hong Kong, Hong Kong, China


Bus emissions - Sweden



- Measurement of primary emissions
- Secondary aerosol production using same oxidation flow reactor (OFR)
- Relevance to HK: majority of franchised buses use similar engine and exhaust treatment technologies

Lindholmen, Gothenburg, Sweden

Bus emissions - Sweden



- Acceleration-phase
- Exhaust-plume probing measurements

Instrument					
CO ₂ NO NOx	Gas Analyzers				
PN	Condensation Particle				
	Counter (CPC)				
PNSD	Engine Exhaust Particle				
	Sizer Spectrometer (EEPS)				
CO, NO, HC	AccuScan RSD 3000,				
VOC, POA, SOA	High-Resolution Time-of-				
	Flight Chemical-Ionization				
	Mass Spectrometer (CIMS)				

Bus emissions - Sweden

Fleet characteristics

EU standard	Engine technology	Fuel	Exhaust after- treatment system	Number of buses	Number of passages	Percentage of plumes with detectable CO ₂ a		
Euro III ^b	Diesel	DSL ^c	SCR+DPF	2	64	88%		
Euro IV	Diesel	RME^d	SCR	10	58	94%		
Euro V	Diesel	DSL ^c	SCR	15	526	87%		
Euro V	Diesel	RME^d	SCR	56	423	95%		
Euro V	Diesel	HVO ^e	SCR	59	432	97%		
Euro V	Diesel	HVO ^e	EGR+DPF	5	26	91%		
Euro V	Diesel _{HEV}	HVO ^e -electric	SCR	26	261	75%		
EEV	CNG	CNG ^f	-	50	533	67%		
Euro VI	Diesel _{HEV}	RME ^d -electric	SCR+EGR+DPF	4	22	63%		
Euro VI	Diesel _{HEV}	HVO ^e -electric	SCR+EGR+DPF	7	350	47%		
TOTAL	-	-	-	<mark>234</mark>	<mark>2695</mark>	-		

^aMinimum threshold


HEV =Hybrid Electric Vehicle

21

^bRetrofitted with SCR (Selective Catalytic Reduction), DPF (Diesel Particulate Filter), EGR = Exhaust Gas Recirculation

^cDSL = diesel fuel; ^d RME = Rapeseed Methyl Ester; ^e HVO = Hydrotreated Vegetable Oil

^fCNG = Compressed Natural Gas: methane, natural and biogas mixture



- ➤ Significant reductions in PM, HC and CO emissions with biofuels compared to diesel
- ➤ Significant increase in NOx emissions due to higher fuel oxygen content in biofuels






- Not fuel dependent

Primary


-fuel dependent




- ➤ The variation of median SA PF was less significant compared to that of MdEF_{PM}
- ➤ The SA PF were high for all fuel types even when the primary emissions were low




Truck emission measurements on a ramp of a freeway in Gothenburg





- ➤ Decreasing EF_{PM} and EF_{PN} with increasingly stringent Euro standards.
- Drastically lower EF_{PM} for Euro VI HDTs but highly scattered EF_{PN}.
- ➤ Average reduction of 13-97% for regulated pollutants from Euro V to Euro VI.





Truck emission measurements: non-regulated and secondary pollutants





- Most of non-regulated gaseous pollutants, e.g. carboxylic acids and byproducts from the urea-SCR exhaust, the average emissions decreased by as high as 90% from Euro V to Euro VI.
- > Euro VI HDTs generate the lowest secondary particle mass compared with other older diesel vehicles.

(In preparation)

Theme 3


Physico-chemical processes in the urban atmosphere (city scale)

Highly polluted environments: haze event in China

- Higher mass fraction of secondary inorganic aerosols and secondary organic aerosols at higher relative humidity: important role of multiphase chemistry during the haze events (Li et al.)
- Missing sulfate production pathways
 - Importance of SO₂ oxidation by NO₂ under high ionic strength conditions (e.g., 60
 M) (Bloss et al.)
 - Potential enhancement of sulfate production by multiphase photochemistry
 (From CKC group)(1-3)
- (1) Gen et al., ES&T Lett., 6(2), 86-91, 2019. (ES&T Lett Best Paper Award)
- (2) Gen et al., ES&T., 53(15), 8757-8766, 2019.
- (3) Zhang et al., ES&T., 54(7), 3831-3839, 2020.

Diel variations of PM1 mass concentrations of the major components under various RH conditions (Li et al.)


The Gucheng site, 100 km southwest of Beijing (11 Nov – 24 Dec 2018)



- Substantial increase in inorganic mass $(SO_4^{2-} \text{ and } NO_3^{-})$ from low RH, to high RH, and to the fog case (RH = 100%)
- Similar but lesser increasing trend of SOA with RH
- Highlighting enhanced formation of secondary species with increasing RH, due to the multiphase processes

Modeled sulfate formation rates via SO_2 oxidation by OH, dissolved H_2O_2 , O_3 , TMI, and NO_2 : importance of NO_2 under high ionic strength (I) conditions (Bloss et al)


Timeseries modeled sulfate formation rates



- Much low sulfate formation rates via SO_2 oxidation by OH, dissolved H_2O_2 , O_3 , TMI under high ionic strength conditions
- Significant increase in the predicted overall formation rate of sulfate due to SO₂ oxidation by NO₂ under high ionic strength conditions, comparable to field observations

$$k_{NO2}(I) = k_{NO2}(I=0) \times 10^b I$$

Proposed mechanism for heterogeneous oxidation of SO₂ into sulfate during nitrate photolysis




- In-particle oxidants produced from nitrate photolysis at > 290 nm irradiation: OH, NO₂, and NO₂⁻ /HNO₂
- 4 pathways for oxidation of dissolved SO₂ by: (1) NO₂, (2) OH,
 (3) NO₂-/HNO₂ (N(III)), (4) H₂O₂
- Organics react with OH to produce O₂-/HO₂
- Self reaction of O₂-/HO₂ for H₂O₂ production


Gen et al. (2019ab)

Methodology: In-situ Raman/Flow cell system to quantify sulfate production

- Sample: NH₄NO₃ (+ halide ions)
- SO_2 : 7.7 ppm
- UV lamp: 250-nm (Mercury) or 300-nm (LED)




Sulfate production during nitrate photolysis at 300nm




- (i) Dark experiment (AN + SO₂ in air) and (ii)
 AN photolysis without SO₂ (AN + UV in air):
 no sulfate production
- NaCl photolysis with SO₂ (NaCl + UV + SO₂ in air): significantly low sulfate production
- AN photolysis with SO₂ in air (AN + UV + SO₂ in air) and N₂ (AN + UV + SO₂ in N₂): sulfate production
- AN photolysis with SO₂ and delay irradiation (AN + delay UV in air): significant sulfate production during irradiation

Nitrate photolysis vs other mechanisms in sulfate production under typical haze conditions in Beijing




- Nitrate photolysis rate ~ 10⁻⁵ M s⁻¹:
 - \checkmark pH > 4, Total sulfate production rate: ~1 μg m⁻³ h⁻¹ \rightarrow close to missing sulfate production rate ~3 μg m⁻³ h⁻¹
- 4.5 < pH < 5.5 (typical pH range under haze event)
 - ✓ Nitrate photolysis process can contribute up to ~60% of sulfate production in aqueous particles.

Halide-induced enhancement of nitrate photolysis leading to increased sulfate production



- High surface propensity of halide ions (Cl⁻, Br⁻ and l⁻).
- Halide ions prefer to reside at the surface → increased counterion (i.e., cation) concentration → attract more nitrate ions to the surface.
- Reduced solvent-cage effect at the interface.

Implementation of the proposed mechanism in air quality model and on-going work



• $\gamma_{SO2} = 1.64 \times j_{NO3} \times [NO_3]$ according to our experimental data

Gen et al., ES&T., 53(15), 8757-8766, 2019.

 Nitrate photolysis pathways can contribute to 15% to 65% to sulfate gap when EF* increase from 10 to 100 in haze events in Beijing

Zheng et al. *ES&T Lett* 7.9 (2020): 632-638.

*EF: enhancement factor of nitrate photolysis rate constant relative to that of gas phase


 Oxidation of Glyoxal during nitrate photolysis under 300-nm

Zhang et al. To be submitted.

Importance of new particle formation (NPF) in air quality in mega cities (Beijing)

- NPF leading to haze episodes (Kulmala et al.)
- Survival of clusters and newly formed particles governed by background aerosols through the coagulation scavenging (Deng et al.)
- Necessity for improved methods for more accurately quantifying the formation and the growth rates of newly formed particles in polluted urban environments (Deng et al.)


Main mechanisms of NPF and subsequent particle growth leading haze episodes (Kulmala et al.)




- NPF initiating almost all present-day haze episodes
- Delaying the buildup of haze episodes by 1–3 days by the reduction in the growth rate of freshly formed particles

Number concentrations of sub-3 nm particles and occurrence frequencies of NPF under different condensation sink (CS)

(Deng et al.)



- At low CS, frequent NPF events and high concentrations of sub-3 nm particles
- At high CS, rarely observed NPF events and the concentrations of below 10³ cm⁻³
- Importance of the coagulation scavenging in governing the survival of clusters and newly formed particles




Vertical profiles: effect of surface emissions in boundary layer

- More pronounced vertical gradient for reactive isoprene and monoterpenes in comparison to less reactive CO, benzene, and toluene (Kim et al.)
- More oxidized compounds in submicron particles at 260 m than at ground level (Xie et al.)
- Little obvious difference in aerosol CCN and hygroscopic behavior at different heights, suggesting the usefulness of surface hygroscopicity measurements (Hu et al.)

Theme 4


Health Effects

PM_{2.5} air quality improvements over China contributed by clean air policies during 2010-2030 (Cheng et al.)



- In 2010, more than 91.5% and 30% of the population exposed above 35 μg/m³, and 75 μg/m³
- Rapid decrease in PM_{2.5} exposures across China with the clean air policies to 26.4 μg/m³ by 2030
- By 2030, the majority of China's population (81.7%, 1143.5 million people) exposed below the National Ambient Air Quality Standard (35 μg/m³)

Contributions of $PM_{2.5}$ component variations from 2000 to 2010 to changes in deaths attributable to $PM_{2.5}$ mixture (Xue et al.)




- According to CS-EMM, BC was the leading contributor to mortality burden
- PRD, **BC** contributed to 6.4%, 6.2%, 5.8%, and 9.0% of changes in total concentration of PM_{2.5}, but to 46.7%, 49.4%, 44.3%, and 51.3% of associated changes in mortality, respectively.

CS-EMM: Components-Specific Exposure-Mortality Model.

BTH: Beijing-Tianjin-Hebei YRD: Yangtze River Delta PRD: Pearl River Delta

Association with health effects: personal exposure vs. ambient measurements (Han et al.)



- Three biomarkers for the inflammatory effects of air pollution: exhaled NO, Monocyte, and CRP
- Among all participants, significant association of personal exposure to $PM_{2.5}$ with an increase in all the three inflammatory biomarkers
- In general, insignificant association of the ambient exposure with the health effects

CRP: C reactive protein

Going forward

Regional scale air quality:

 Rich literature: emissions, source apportionment, ambient characterization, SOA, BC/OC, SNA, NPF and growth, ammonia, pH, CCN

Health:

- Aging and dense population require better air quality
- Personal exposure assessment vs ambient concentrations
- BC

Indoor and neighborhood scale air quality studies:

- Emission inventories: traffic including non tail pipe emissions, cooking emissions
- Modeling including the above emissions, NPF and growth, and atmospheric chemistry and SOA formation
- Measurements

COVID Lockdown:

- Interesting "new" atmospheric chemistry and physics
- Increased importance of secondary processes
- Projection of the emissions, air quality, and personal exposure under the New Normal and Green Recovery

Air pollution in megacities

- LA, London, Beijing, Shanghai, Mexico City, Delhi, Bangkok, Paris, Seoul etc...

CRITICAL REVIEW

ISSN 1047-3289 J. Air & Waste Manage. Assoc. 54:644-680 Copyright 2004 Air & Waste Management Association



Mario J. Molina



Luisa T. Molina

Megacities and Atmospheric Pollution

Mario J. Molina and Luisa T. Molina

Massachusetts Institute of Technology, Cambridge,

Massachusetts





Atmospheric Environment 42 (2008) 1-42

ATMOSPHERIC ENVIRONMENT

www.elsevier.com/locate/atmosenv

Review

Air pollution in mega cities in China

Chak K. Chan^{a,*}, Xiaohong Yao^{a,b}

^aDepartment of Chemical Engineering, Hong Kong University of Science and Technology, Clear Water Bay, Hong Kong, China ^bKey Laboratory of Marine Environmental Science and Ecology, Ministry of Education, Ocean University of China, 238 Songling Road, Oingdao 266100, PR China

Received 10 May 2007; received in revised form 4 September 2007; accepted 4 September 2007


Acknowledgments

Funding Support

Environment & Conservation Fund



University Grants Committee & Research Grants Council



National Science Foundation of China



N S F National Natura Foundation of C

CKC research group

Dr. Berto Lee (CE), Dr. Yongjie Li (U. Macao), and Dr. DanDan Huang (SAES) (AMS measurements in MK)

Dr. Teng Yu Liu (U. Toronto) (Cooking SOA)

Dr. Qianyun Liu and Ms Liyuan Zhou (Bus measurements in Sweden)

Dr. Masao Gen (Kanazawa U.) and Ms Ruifeng Zhang (Sulfate)

Collaborators

Prof. Guo Hai – Hong Kong Polytechnic University



Prof. Jianzhen Yu — Hong Kong University of Science and Technology



Prof. Man Nin Chan – Chinese University of Hong Kong



Prof. Mattias Hallquist – University of Gothenburg



Åsa Hallquist - IVL Swedish Environmental Research Institute



Prof. Wang Xinming - Guangzhou Institute of Geochemistry, CAS



Air Science Group – Environmental Protection Department, HKEPD

